

ON INTRINSIC INVARIANCE IN GURZADYAN-XUE COSMOLOGICAL MODELS

H. KHACHATRYAN, G.V. VERESHCHAGIN AND G. YEGORIAN

*ICRANet, P.le della Repubblica 10, 65100 Pescara, Italy and
ICRA, Dip. Fisica, Univ. "La Sapienza", P.le A. Moro 5, 00185 Rome, Italy
E-mail: veresh@icra.it*

Analysis of cosmological solutions of GX dark energy models shows the presence of a separatrix in the phase space of solutions which divides them into two classes: Friedmannian-like with initial singularity and non-Friedmannian solutions. The invariants are found, that reveal the intrinsic symmetry in GX models.

The formula for dark energy ¹

$$\rho_{GX} = \frac{\pi c^4}{8 G} \frac{1}{a^2}, \quad (1)$$

c , G and a are the speed of light, gravitational constant, and scale factor of the Universe, respectively, derived by Gurzadyan and Xue predicts the value of dark energy density in agreement with present observations ². This formula leads to a possibility of variation of physical constants such as speed of light, gravitational constant, cosmological term. Classification of models based on the GX scaling (1) is given in ³. Analysis of the cosmological equations for GX models is performed in ⁴.

It was shown that each GX model is characterized by the critical value of the density parameter $\Omega_m \approx 2/3$ which represents a separatrix in the space of solutions of cosmological equations ⁵. To illustrate the presence of separatrix we provide in fig.1 phase portraits for expansion stage with zero spatial curvature. Numeration of models is as follows: I - variable dark energy density, II - variable speed of light, constant cosmological term, III - variable gravitational constant, IV - variable speed of light, constant dark energy density (details see e.g. in ⁴).

Phase space variables are

$$\alpha = \frac{2}{\pi} \arctg(a), \quad m = \frac{2}{\pi} \arctg(\mu), \quad (2)$$

where μ is the matter density.

The separatrix divides the space of solutions into two classes: Friedmannian-like with initial singularity and non-Friedmannian solutions starting with vanishing




Figure 1. Phase portraits for models I-IV. Directions of phase trajectories are shown by arrows. See detailed explanations in the text.

density⁵. The value of separatrix is given by

$$\Omega_m = \frac{2}{3} \frac{1}{(1 - k/\pi^2)}. \quad (3)$$

The upper left corner in each diagram at fig.1 corresponds to the Friedmann cosmological singularity with $\mu \rightarrow \infty$ and $a \rightarrow 0$. In the lower right corner $\mu \rightarrow 0$ and $a \rightarrow \infty$. The separatrix in each diagram is shown by a thick curve. Above the separatrix solutions start with a classical singularity and end up with infinite scale factor and zero density (or constant density for models II and III) but the solutions below the separatrix start at a positive scale factor and zero density and tend to

the same limit.

The presence of separatrix in GX models is a crucial difference with the Friedmann solutions within General Relativity. It is shown that such symmetries are due to existence of invariant in GX models⁶; the value of separatrix can be obtained from expression of vanishing invariant.

The continuity equation for GX models reads³

$$\dot{\mu} + 3\frac{\dot{a}}{a}\mu = -\dot{\mu}_{GX} + (\mu + \mu_{GX})\left(2\frac{\dot{c}}{c} - \frac{\dot{G}}{G}\right), \quad (4)$$

where $\mu_{GX} \equiv \rho_{GX}/c^2$ is dark energy mass density. Invariant was found for this equation⁶ which has the form

$$\frac{\mu a^3 G}{c^2} - \frac{\pi a}{4} = const \equiv b_m. \quad (5)$$

The value $b_m = 0$ corresponds exactly to the above condition (3).

Apart from the demonstrated symmetry, present in all GX models, which is linked with the form of (1), it is shown that GX models agree well with supernovae and radio galaxies data⁷. Perturbation dynamics analysis⁸ also shows some interesting features of GX models, in particular that density fluctuations grow also during curvature- or vacuum-dominated stages, following matter-dominated stage of expansion, unlike perturbations within Friedmannian models.

References

1. V.G. Gurzadyan, S.-S. Xue in: "From Integrable Models to Gauge Theories; volume in honor of Sergei Matinyan", ed. V.G. Gurzadyan, A.G. Sedrakian, p.177, *World Scientific*, 2002; *Mod. Phys. Lett.* **A18** (2003) 561 [astro-ph/0105245], see also [astro-ph/0510459].
2. S.G. Djorgovski, V.G. Gurzadyan, Nucl.Phys. B (in press), *astro-ph/0610204*.
3. G.V. Vereshchagin, *Mod. Phys. Lett.* **A21** (2006) 729.
4. G.V. Vereshchagin and G. Yegorian, *Class. Quantum Grav.* **23** (2006) 5049.
5. G.V. Vereshchagin and G. Yegorian, *Phys. Lett.* **B636** (2006) 150.
6. H.G. Khachatryan, *astro-ph/0608649*.
7. G.V. Vereshchagin and G. Yegorian, *astro-ph/0604566*.
8. G.V. Vereshchagin and G. Yegorian, *astro-ph/0610197*; see also this volume.